arXiv:0707.3909v1 [math-ph] 26 Jul 2007

Schrödinger operators on armchair nanotubes. I

Andrey Badanin [*] Jochen Brüning [†] Evgeny Korotyaev [‡]
Igor Lobanov [§]

February 5, 2008

Abstract

We consider the Schrödinger operator with a periodic potential on quasi-1D models of armchair single-wall nanotubes. The spectrum of this operator consists of an absolutely continuous part (intervals separated by gaps) plus an infinite number of eigenvalues with infinite multiplicity. We describe all eigenfunctions with the same eigenvalue. We define a Lyapunov function, which is analytic on some Riemann surface. On each sheet, the Lyapunov function has the same properties as in the scalar case, but it has branch points, which we call resonances. In example we show the existence of real and complex resonances for some specific potentials.

1 Introduction and main results

Consider the Schrödinger operator $\mathscr{H} = -\Delta + \mathscr{V}_q$ with a periodic potential $\mathscr{V}_q$ on so called armchair graph $\Gamma^N, N \geqslant 1$. In order to describe the graph Γ^N, we introduce the fundamental cell $\widetilde{\Gamma} = \cup_{j\in\mathbb{N}_6}\widetilde{\Gamma}_j \subset \mathbb{R}^2$, where $\widetilde{\Gamma}_j = \{x = \widetilde{\mathbf{r}}_j + t\mathbf{e}_j, t \in [0,1]\}$ are edges of length 1,

$$\mathbb{N}_m = \{1,2,..,m\}, \quad \mathbf{e}_1 = \mathbf{e}_6 = \frac{1}{2}(1,\sqrt{3}), \quad \mathbf{e}_2 = \mathbf{e}_4 = (1,0), \quad \mathbf{e}_3 = -\mathbf{e}_5 = \frac{1}{2}(1,-\sqrt{3}),$$
$$\widetilde{\mathbf{r}}_1 = (0,0), \quad \widetilde{\mathbf{r}}_2 = \widetilde{\mathbf{r}}_5 = \widetilde{\mathbf{r}}_1 + \mathbf{e}_1, \quad \widetilde{\mathbf{r}}_3 = \widetilde{\mathbf{r}}_6 = \widetilde{\mathbf{r}}_2 + \mathbf{e}_2, \quad \widetilde{\mathbf{r}}_4 = \widetilde{\mathbf{r}}_3 + \mathbf{e}_3. \quad (1.1)$$

We define the strip graph $\widetilde{\Gamma}^N$ by

$$\widetilde{\Gamma}^N = \cup_{(n,k)\in\mathbb{Z}\times\mathbb{N}_N}(\widetilde{\Gamma} + k\mathbf{e}_h + n\mathbf{e}_v) \subset \mathbb{R}^2, \quad \mathbf{e}_h = (3,0), \quad \mathbf{e}_v = (0,\sqrt{3}).$$

---

[*] Arkhangelsk State Technical University, e-mail: a.badanin@agtu.ru

[†] Institut für Mathematik, Humboldt Universität zu Berlin, e-mail: bruening@math.hu-berlin.de

[‡] Correspondence author. Institut für Mathematik, Humboldt Universität zu Berlin, Rudower Chaussee 25, 12489, Berlin, Germany, e-mail: evgeny@math.hu-berlin.de

[§] Mathematics Department, Saint-Petersburg State University of Information Technologies, Mechanics and Optics, 1971101, Saint-Petersburg, Sablinskaya 14, email: lobanov@mathdep.ifmo.ru



Figure 1: Armchair graph for $N = 10$ and for $N = 1$.

Vertices of $\widetilde{\Gamma}^N$ are given by $\widetilde{\mathbf{r}}_j + k\mathbf{e}_h + n\mathbf{e}_v, (n, j, k) \in \mathbb{Z} \times \mathbb{N}_6 \times \mathbb{N}_N$. If we identify the vertices $\widetilde{\mathbf{r}}_1 + n\mathbf{e}_v$ and $\widetilde{\mathbf{r}}_1 + N\mathbf{e}_h + n\mathbf{e}_v$ of $\widetilde{\Gamma}^N$ for each $n \in \mathbb{Z}$, then we obtain the graph Γ^N by

$$\Gamma^N = \cup_{\omega \in \mathcal{Z}} \Gamma_\omega, \quad \omega = (n, j, k) \in \mathcal{Z} = \mathbb{Z} \times \mathbb{N}_6 \times \mathbb{Z}_N, \quad \mathbb{Z}_N = \mathbb{Z}/(N\mathbb{Z}),$$

where $\Gamma_\omega = \widetilde{\Gamma}_j + k\mathbf{e}_h + n\mathbf{e}_v$, see Fig. 1, 2. Let $\mathbf{r}_\omega = \widetilde{\mathbf{r}}_j + k\mathbf{e}_h + n\mathbf{e}_v$ be a starting point of the edge Γ_ω. We have the coordinate $x = \mathbf{r}_\omega + t\mathbf{e}_j$ and the local coordinate $t \in [0, 1]$ on Γ_ω. Thus we give an orientation on the edge. For each function y on Γ^N we define a function $y_\omega = y|_{\Gamma_\omega}, \omega \in \mathcal{Z}$. We identify each function y_ω on Γ_ω with a function on $[0, 1]$ by using the local coordinate $t \in [0, 1]$. Define the Hilbert space $L^2(\Gamma^N) = \oplus_{\omega \in \mathcal{Z}} L^2(\Gamma_\omega)$. Let $C(\Gamma^N)$ be the space of continuous functions on Γ^N. We define the Sobolev space $W^2(\Gamma^N)$ that consists of all functions $y = (y_\omega)_{\omega \in \mathcal{Z}} \in L^2(\Gamma^N), (y''_\omega)_{\omega \in \mathcal{Z}} \in L^2(\Gamma^N)$ and satisfy
Kirchhoff Boundary Conditions: $y \in C(\Gamma^N)$ *and for each vertex* A *of* Γ^N *satisfies:*

$$\sum_{\omega \in E_A} (-1)^b y'_\omega(b) = 0, \quad \textit{where} \quad E_A = \{\omega \in \mathcal{Z} : A \in \Gamma_\omega\}, \quad b = b(\omega, A), \tag{1.2}$$

where if $A = \mathbf{r}_\omega$ *is a starting point of* Γ_ω *(i.e.* $t = 0$ *at* A*), then* $b(\omega, A) = 0$,
if $A = \mathbf{r}_\omega + \mathbf{e_j}$ *is an endpoint of* Γ_ω *(i.e.* $t = 1$ *at* A*), then* $b(\omega, A) = 1$.

The Kirchhoff Conditions (1.2) mean that the sum of derivatives of y at each vertex of Γ^N equals 0 and the orientation of edges gives the sign ±. Our operator $\mathscr{H}$ on Γ^N acts in the Hilbert space $L^2(\Gamma^N)$ and is given by $(\mathscr{H} y)_\omega = -y''_\omega + q y_\omega$, where $y = (y_\omega)_{\omega \in \mathcal{Z}} \in \mathfrak{D}(\mathscr{H}) = W^2(\Gamma^N)$ and $(\mathscr{V}_q y)_\omega = q y_\omega, q \in L^2(0, 1)$. Note that the orientation of edges is not important



Figure 2: A piece of a nanotube Γ^N. The fundamental domain is marked by a bold line.

for the case of even potentials $q \in L^2_{even}(0,1) = \{q \in L^2(0,1) : q(t) = q(1-t), t \in (0,1)\}$. The standard arguments (see [KL]) yield that $\mathscr{H}$ is self-adjoint.

For the convenience of the reader we briefly describe the structure of carbon nanotubes, see [Ha], [SDD]. Graphene is a single 2D layer of graphite forming a honeycomb lattice, see Fig. 3. A carbon nanotube is a honeycomb lattice "rolled up" into a cylinder, see Fig. 1. In carbon nanotubes, the graphene sheet is "rolled up" in such a way that the so-called chiral vector $\Omega = N_1\Omega_1 + N_2\Omega_2$ becomes the circumference of the tube, where Ω_1, Ω_2 are defined in Fig 3. The chiral vector Ω, which is usually denoted by the pair of integers (N_1, N_2), uniquely defines a particular tube. Tubes of type $(N, 0)$ are called zigzag tubes. (N, N)-tubes are called armchair tubes.

Our Schrödinger operator can be considered as a model Hamiltonain for π-electrons in armchair single-wall carbon nanotubes (SWCN) which attract tremendous interest from fundamental science and technological perspectives [AL]. The best known model of SWCN is a discrete one obtained in the nearest-neighborhood tight-binding approximation proposed in 1992 (see the articles [HSO], [MDW], [SFDD] and review in [SDD]) shortly after the discovery of carbon nanotubes [Ii]. Recently, the model was applied to SWCN (in particular to armchair configuration) subjected to an electric field [PBRL]. Being qualitative satisfactory, the tight binding approximation leads sometimes to quantitative errors, which are commonly addressed to curvature effects, see [OHKL]. However, as was shown in [RMT], the tight-binding including up to third-nearest neighbors significantly improves approximation. To turn on interaction between all atoms in nanotube, one can proceed to the model considered in [AT], where carbon atoms was modelled by zero-range potentials on a cylinder. In the present article we use an intermediate model between the mentioned discrete and



Figure 3: The honeycomb lattice of nanotube. The unit cell is spanned by the vectors Ω_1 and Ω_2. The type of the nanotube is defined by the pair $(N_1, N_2) \in \mathbb{N}^2$, $N_1 \geqslant N_2$, and corresponding chiral vector $\Omega = N_1\Omega_1 + N_2\Omega_2$.

two-dimensional ones, namely we deal with quantum network models which attract lot of attention over the last decade, see [Ku]. The considered model was introduced by Pauling [Pa] and was systematically developed in the series of articles by Ruedenberg and Scherr [RS]. Further progress had been made toward periodic systems by Coulson in [C] where a network model of graphite layer was worked out. A network model of a crystal with non-trivial potentials along bonds was studied by Montroll [M]. Alexander [A] noticed simple relations between spectra of quantum graphs and combinatorial properties of discrete ones which became common for spectral analysis on quantum networks at the present time.

The simplest pure carbon nanotube consists of atoms with covalent bond, when roughly speaking the potential of electric field between two atoms is even. The situation is more complicated, for example, in the model of a nitrogen-carbon-boron nanotube (see [Ha], Ch.7.3), where the bond between atoms is ionic and the corresponding potential is non-even.

Recall the needed properties of the Hill operator $\widetilde{H}y = -y'' + q(t)y$ on the real line with a periodic potential $q(t+1) = q(t), t \in \mathbb{R}$. The spectrum of $\widetilde{H}$ is purely absolutely continuous and consists of intervals $\widetilde{\sigma}_n = [\widetilde{\lambda}_{n-1}^+, \widetilde{\lambda}_n^-], n \geqslant 1$. These intervals are separated by the gaps $\widetilde{\gamma}_n = (\widetilde{\lambda}_n^-, \widetilde{\lambda}_n^+)$ of length $|\widetilde{\gamma}_n| \geqslant 0$. If a gap $\widetilde{\gamma}_n$ is degenerate, i.e. $|\widetilde{\gamma}_n| = 0$, then the corresponding segments $\widetilde{\sigma}_n, \widetilde{\sigma}_{n+1}$ merge. For the equation $-y'' + q(t)y = \lambda y$ on the real line we define the fundamental solutions $\vartheta(t, \lambda)$ and $\varphi(t, \lambda), t \in \mathbb{R}$ satisfying $\vartheta(0, \lambda) = \varphi'(0, \lambda) = 1, \vartheta'(0, \lambda) = \varphi(0, \lambda) = 0$. Let $\varphi_1 = \varphi(1, \cdot), \vartheta_1 = \vartheta(1, \cdot), \varphi_1' = \varphi'(1, \cdot), \vartheta_1' = \vartheta'(1, \cdot)$. The corresponding monodromy matrix $\widetilde{\mathcal{M}}$, the Lyapunov function F, and the standard function F_- are given by

$$\widetilde{\mathcal{M}} = \begin{pmatrix} \vartheta_1 & \varphi_1 \\ \vartheta_1' & \varphi_1' \end{pmatrix}, \quad F = \frac{\varphi_1' + \vartheta_1}{2}, \quad F_- = \frac{\varphi_1' - \vartheta_1}{2}. \tag{1.3}$$

The sequence $\widetilde{\lambda}_0^+ < \widetilde{\lambda}_1^- \leqslant \widetilde{\lambda}_1^+ < ...$ is the spectrum of the equation $-y'' + qy = \lambda y$ with two periodic boundary conditions, that is $y(t+2)=y(t), t\in\mathbb{R}$. Here equality $\widetilde{\lambda}_n^-=\widetilde{\lambda}_n^+$ means that $\widetilde{\lambda}_n^\pm$ is an eigenvalue of multiplicity 2. Note that $F(\widetilde{\lambda}_n^\pm)=(-1)^n,\ n\geqslant 1$. The lowest eigenvalue $\widetilde{\lambda}_0^+$ is simple, $F(\widetilde{\lambda}_0^+)=1$, and the corresponding eigenfunction has period 1. The eigenfunctions corresponding to $\widetilde{\lambda}_n^\pm$ have period 1 if n is even, and they are anti-periodic, that is $y(t+1)=-y(t),\ t\in\mathbb{R}$, if n is odd. The derivative of the Lyapunov function has a zero $\widetilde{\lambda}_n$ in each interval $[\lambda_n^-,\lambda_n^+]$, that is $F'(\widetilde{\lambda}_n)=0$. Let $\mu_n, n\geqslant 1$, be the spectrum of the problem $-y''+qy=\lambda y, y(0)=y(1)=0$ (the Dirichlet spectrum), and let $\nu_n, n\geqslant 0$, be the spectrum of the problem $-y''+qy=\lambda y, y'(0)=y'(1)=0$ (the Neumann spectrum). Define the set $\sigma_D=\{\mu_n, n\geqslant 1\}$ and note that $\sigma_D=\{\lambda\in\mathbb{C}:\varphi(1,\lambda)=0\}$. It is well-known that $\mu_n,\nu_n\in[\widetilde{\lambda}_n^-,\widetilde{\lambda}_n^+], n\geqslant 1$, and $\nu_0\leqslant\widetilde{\lambda}_0^+$.

For simplicity we shall denote $\Gamma_{\alpha,1}\subset\Gamma^1$ by Γ_α, for $\alpha=(n,j)\in\mathcal{Z}_1=\mathbb{Z}\times\mathbb{N}_6$. Thus $\Gamma^1=\cup_{\alpha\in\mathcal{Z}_1}\Gamma_\alpha$, see Fig 1. In Theorem 1.1 we will show that $\mathscr{H}$ is unitarily equivalent to $H=\oplus_1^N H_k$, where the operator H_k acts in the Hilbert space $L^2(\Gamma^1)$ and is given by $(H_k f)_\alpha=-f''_\alpha+qf_\alpha,\ (f_\alpha)_{\alpha\in\mathcal{Z}_1},(f''_\alpha)_{\alpha\in\mathcal{Z}_1}\in L^2(\Gamma^1)$, and the components $f_\alpha,\alpha\in\mathcal{Z}_1$ satisfy **the Kirchhoff conditions**:

$$\begin{aligned}&f_{n,1}(1)=f_{n,2}(0)=f_{n,5}(0),\qquad f_{n,2}(1)=f_{n,3}(0)=f_{n,6}(0),\\ &\qquad f_{n,3}(1)=f_{n,4}(0)=f_{n-1,6}(1),\quad s^k f_{n,4}(1)=f_{n,1}(0)=f_{n-1,5}(1),\quad s=e^{i\frac{2\pi}{N}},\end{aligned}\tag{1.4}$$

$$\begin{aligned}&f'_{n,1}(1)-f'_{n,2}(0)-f'_{n,5}(0)=0,\quad f'_{n,2}(1)-f'_{n,3}(0)-f'_{n,6}(0)=0,\\ &\qquad f'_{n,3}(1)-f'_{n,4}(0)+f'_{n-1,6}(1)=0,\quad s^k f'_{n,4}(1)-f'_{n,1}(0)+f'_{n-1,5}(1)=0.\end{aligned}\tag{1.5}$$

Our analysis is close to previous papers [KL], [KL1] about the Schrödinger operators on the zigzag graphs. We also introduce the monodromy matrix and the Lyapunov functions. We study the properties of these functions and here we essentially use the results and techniques from the papers [BBK],[CK], [K1] and [KL].

For the operator H_k we define fundamental solutions $\vartheta_k^{(\nu)}=(\vartheta_{k,\alpha}^{(\nu)})_{\alpha\in\mathcal{Z}_1},\varphi_k^{(\nu)}=(\varphi_{k,\alpha}^{(\nu)})_{\alpha\in\mathcal{Z}_1}$, $\nu=1,2$ which satisfy

$$\begin{aligned}&-\Delta y+\mathscr{V}_q y=\lambda y,\quad on\ \ \Gamma^1,\qquad \text{the Kirchhoff Boundary Conditions (1.4),(1.5)}\\ &\qquad \Theta_{k,-1}(1,\lambda)=\Phi'_{k,-1}(1,\lambda)=I_2,\quad \Theta'_{k,-1}(1,\lambda)=\Phi_{k,-1}(1,\lambda)=0,\end{aligned}\tag{1.6}$$

where $I_n, n\geqslant 2$ is the $n\times n$ identity matrix and

$$\Theta_{k,n}=\begin{pmatrix}\vartheta_{k,n,5}^{(1)} & \vartheta_{k,n,5}^{(2)}\\ \vartheta_{k,n,6}^{(1)} & \vartheta_{k,n,6}^{(2)}\end{pmatrix},\quad \Phi_{k,n}=\begin{pmatrix}\varphi_{k,n,5}^{(1)} & \varphi_{k,n,5}^{(2)}\\ \varphi_{k,n,6}^{(1)} & \varphi_{k,n,6}^{(2)}\end{pmatrix}.\tag{1.7}$$

We define the monodromy matrix (determined by the fundamental solutions on Γ_0^1) by

$$\mathcal{M}_k(\lambda)=\begin{pmatrix}\Theta_{k,0} & \Phi_{k,0}\\ \Theta'_{k,0} & \Phi'_{k,0}\end{pmatrix}(1,\lambda).\tag{1.8}$$

We formulate our first result about the monodromy matrix.



Figure 4: The supports of eigenfunctions: a) $\psi^{(0,1)}$, b) $\psi^{(0,2)}$.

Theorem 1.1. *i) The operator* $\mathscr{H}$ *is unitarily equivalent to* $H = \oplus_1^N H_k$.
ii) Let $k \in \mathbb{Z}_N$. *Then for any* $\lambda \in \mathbb{C} \setminus \sigma_D$ *there exist unique fundamental solutions* $\vartheta_k^{(\nu)} = (\vartheta_{k,\alpha}^{(\nu)})_{\alpha \in \mathcal{Z}_1}, \varphi_k^{(\nu)} = (\varphi_{k,\alpha}^{(\nu)})_{\alpha \in \mathcal{Z}_1}, \nu = 1, 2$. *Moreover, each of the functions* $\vartheta_{k,\alpha}^{(\nu)}(x, \lambda)$, $\varphi_{k,\alpha}^{(\nu)}(x, \lambda)$, $x \in \Gamma^1$ *is analytic in* $\lambda \in \mathbb{C} \setminus \sigma_D$ *and the monodromy matrix* $\mathcal{M}_k(\lambda)$ *satisfies*

$$\det \mathcal{M}_k = 1, \quad \operatorname{Tr} \mathcal{M}_0 = 2(9F^2 - F_-^2 - 1), \quad \operatorname{Tr} \mathcal{M}_k = \operatorname{Tr} \mathcal{M}_0 - 4s_k^2, \quad s_k = \sin \frac{\pi k}{N}, \tag{1.9}$$

$$\operatorname{Tr} \mathcal{M}_0^2 = 72F^2 + \frac{1}{2}(\operatorname{Tr} \mathcal{M}_0)^2 - 4, \quad \operatorname{Tr} \mathcal{M}_k^2 = \operatorname{Tr} \mathcal{M}_0^2 - 8s_k^2 \operatorname{Tr} \mathcal{M}_0 - 4s_{2k}^2, \tag{1.10}$$

$$\mathcal{M}_k(\lambda)^\top J \mathcal{M}_k(\lambda) = J, \qquad \text{where} \quad J = \begin{pmatrix} 0 & \mathbf{j}_2 \\ -\mathbf{j}_2 & 0 \end{pmatrix}, \ \ \mathbf{j}_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \tag{1.11}$$

and the matrix-valued function $\mathcal{R}\mathcal{M}_k\mathcal{R}^{-1}$ *is entire, where* $\mathcal{R} = I_2 \oplus \varphi_1 I_2$.

The monodromy matrix $\mathcal{M}_k$ has poles at the points $\lambda \in \sigma_D$, which are eigenvalues of H_k (see Theorem 1.2), similar to the zigzag tube [KL], [MV].

Define the subspace $\mathcal{H}_k(\lambda) = \{\psi \in \mathfrak{D}(H_k) : H_k\psi = \lambda\psi\}$ for $(\lambda, k) \in \sigma_{pp}(H_k) \times \mathbb{Z}_N$. If $\dim \mathcal{H}_k(\lambda_0) = \infty$ for some $\lambda_0 \in \sigma_{pp}(H_k)$, then we say that $\{\lambda_0\}$ is a flat band. In Theorem 1.2 we describe all flat bands and corresponding eigenfunctions (see Fig. 4).

Theorem 1.2. *Let* $(\lambda, k) \in \sigma_D \times \mathbb{Z}_N$. *Then*
i) Every eigenfunction of $\mathcal{H}_k(\lambda)$ *vanishes at all verteces of* Γ^1.

ii) There exist two functions $\psi^{(0,\nu)} = (\psi^{(0,\nu)}_{\alpha})_{\alpha\in\mathcal{Z}_1} \in \mathcal{H}_k(\lambda), \nu = 1,2,$ *on* Γ^1 *such that* $\operatorname{supp}\psi^{(0,\nu)} \subset \cup_{j=1}^{6}\Gamma_{0,j}$*, each function* $\psi^{(n,\nu)} = (\psi^{(0,\nu)}_{n-m,j})_{(m,j)\in\mathcal{Z}_1} \in \mathcal{H}_k(\lambda), n \in \mathbb{Z}$*. Moreover, each* $f \in \mathcal{H}_k(\lambda)$ *satisfies:*

$$f = \sum_{n\in\mathbb{Z}}(\widehat{f}_{n,1}\psi^{(n,1)} + \widehat{f}_{n,2}\psi^{(n,2)}), \qquad (\widehat{f}_{n,1}, \widehat{f}_{n,2})_{n\in\mathbb{Z}} \in \ell^2 \oplus \ell^2, \tag{1.12}$$

where

$$\widehat{f}_{n,1} = -\varphi_1' f_{n,1}'(0), \quad \widehat{f}_{n,2} = f_{n,2}'(0), \quad \text{if} \quad \varphi_1'^2 = 1, k = 0,$$

$$\widehat{f}_{n,1} = \frac{\varkappa_1 f_{n,6}'(0) + f_{n,3}'(0)}{\varkappa_1\varkappa_2 - 1}, \quad \widehat{f}_{n,2} = \frac{f_{n,6}'(0) + \varkappa_2 f_{n,3}'(0)}{\varkappa_1\varkappa_2 - 1}, \quad \text{if}\, \varphi_1'^2 \neq 1 \text{ or } \varphi_1'^2 = 1, k \neq 0, \tag{1.13}$$

$\varkappa_1 = 1 - s^k\varphi_1'^2, \varkappa_2 = 1 - s^k\varphi_1'^4$*. Moreover, the mapping* $f \to (\widehat{f}_{n,1}, \widehat{f}_{n,2})_{n\in\mathbb{Z}}$ *is a linear isomorphism between* $\mathcal{H}_k(\lambda)$ *and* $\ell^2 \oplus \ell^2$*.*

Using (1.11) we deduce that eigenvalues of $\mathcal{M}_k, k \in \mathbb{Z}_N$ have the form $\tau_{1,\nu}^{\pm 1}, \tau_{2,\nu}^{\pm 1}$.

Theorem 1.3. *Let* $\tau_{k,1}^{\pm 1}, \tau_{k,2}^{\pm 1}$ *be eigenvalues of* $\mathcal{M}_k$ *for some* $k \in \mathbb{Z}_N$*. Then*
(i) The Lyapunov functions $F_{k,\nu} = \frac{1}{2}(\tau_{k,\nu} + \frac{1}{\tau_{k,\nu}}), \nu = 1,2$ *satisfy*

$$F_{k,\nu} = \xi_k - (-1)^{\nu}\sqrt{\rho_k}, \quad \xi_k = \frac{9F^2 - F_-^2 - 1}{2} - s_k^2, \quad \rho_k = (9F^2 - s_k^2)c_k^2 + s_k^2 F_-^2, \tag{1.14}$$

$$D_k(\tau, \lambda) \equiv \det(\mathcal{M}_k(\lambda) - \tau I_4) = (\tau^2 - 2F_{k,1}(\lambda)\tau + 1)(\tau^2 - 2F_{k,2}(\lambda)\tau + 1), \tag{1.15}$$

where $F_{k,\nu}$ *are branches of* $F_k = \xi_k + \sqrt{\rho_k}$ *on a two sheeted Riemann surface* $\mathfrak{R}_k$ *defined by* $\sqrt{\rho_k}$*.*
(ii) If $F_k(\lambda) \in (-1,1)$ *for some* $\lambda \in \mathbb{R}$ *and* λ *is not a branch point of* F_k*, then* $F_k'(\lambda) \neq 0$*.*
(iii) The following identities hold:

$$\sigma(H_k) = \sigma_\infty(H_k) \cup \sigma_{ac}(H_k), \quad \sigma_\infty(H_k) = \sigma_D,$$
$$\sigma_{ac}(H_k) = \{\lambda \in \mathbb{R} : F_{k,\nu}(\lambda) \in [-1,1] \text{ for some } \nu \in \mathbb{N}_2\}. \tag{1.16}$$

(iv) If $q \in L^2_{even}(0,1)$*, then* $\sigma_{ac}(H) = \sigma_{ac}(H_0) = \sigma(\widetilde{H})$*.*

Definition 1. *A zero of* $\rho_k, k \in \mathbb{Z}_N$ *is called a* **resonance** *of* H_k*.*

Let $r_{k,n}^{\pm}, n \geqslant 1$ be zeros of ρ_k and let their labeling be given by $\operatorname{Re} r_{k,1}^{-} \leqslant \operatorname{Re} r_{k,1}^{+} \leqslant \operatorname{Re} r_{k,2}^{-} \leqslant \operatorname{Re} r_{k,2}^{+} \leqslant ...$ We show the existence of real and non-real resonances.

Proposition 1.4. *Let* $q = q_\varepsilon = \frac{1}{v}\delta(t - \frac{1}{2} - c_k\varepsilon - \varepsilon^2), \varepsilon \neq 0, t \in [0,1]$ *for some* $k \notin \{0, \frac{N}{2}\}$*, and let* $n_0 > 1$*. Then for each* $0 < n \leqslant n_0$ *there exist functions* $r_{k,n}^{\pm}(z^2)$*, analytic in the disk* $\{|z| < \varepsilon_n\}$ *with some* $\varepsilon_n > 0$*, such that* $\rho_k(r_{k,n}^{\pm}(\varepsilon), q_\varepsilon) = 0$ *for all* $\varepsilon \in (-\varepsilon_n, \varepsilon_n) \setminus \{0\}$ *and*

$$r_{k,n}^{\pm}(0) = (\pi n)^2, \qquad r_{k,n}^{\pm}(\varepsilon) = (\pi n)^2 - 2\pi n\varepsilon \pm i\frac{4\sqrt{2}(\pi n)^2 s_k}{3\sqrt{c_k}}\varepsilon^{3/2} + O(\varepsilon^2) \quad \text{as} \quad v \to 0. \tag{1.17}$$

Moreover, $\omega_{k,n}^{-}(\varepsilon) = (r_{k,n}^{-}(\varepsilon), r_{k,n}^{+}(\varepsilon)) \subset \mathbb{R}$ *and* $\rho_k(\lambda, q_\varepsilon) < 0$ *for all* $\lambda \in \omega_{k,n}^{-}(\varepsilon), \varepsilon \in (-\varepsilon_n, 0)$*.*



Figure 5: Graph of the Lyapunov function $F_k(\lambda, q_\varepsilon)$ for fixed $\varepsilon \in (0, \varepsilon_1)$. The spectral bands of $H_k(q_\varepsilon)$ are marked by bold lines, counted with multiplicity. Here $E^{k,\pm}_{\nu,n}$ are endpoints of the bands: $E^{k,\pm}_{1,3} = r^{\pm}_{k,2}$ and all other points $E^{k,\pm}_{\nu,n}$ satisfy $F_{k,\nu}(E^{k,\pm}_{\nu,n}) = (-1)^n$, $E^{k,\pm}_{\nu,2n}$ are periodic eigenvalues, $E^{k,\pm}_{\nu,2n+1}$ are antiperiodic eigenvalues.

Remark. (i) Graph of the function $F_k, k \neq \{0, \frac{N}{2}\}$ for $q = q_\varepsilon, \varepsilon \in (0, \varepsilon_1)$ is given by Fig.5.
(ii) The potential $q = q_\varepsilon = \frac{1}{v}\delta(t - \frac{1}{2} - c_k\varepsilon - \varepsilon^2), v \neq 0$ in Proposition 1.4 is not even.
(iii) If $\varepsilon \in (-\varepsilon_n, 0)$, then $F_{k,\nu}(\lambda) \notin \mathbb{R}$ for all $\lambda \in \omega^-_{k,n}(\varepsilon)$, which yields $\omega^-_{k,n}(\varepsilon) \cap \sigma(H_k) = \emptyset$.

There are a lot of papers about the spectral analysis of the Schrödinger operator on periodic graph and periodic nanotubes. Molchanov and Vainberg [MV] consider Schrödinger operators with $q = 0$ on so-called necklace periodic graphs. Korotyaev and Lobanov [KL], [KL1] consider the Schrödinger operator on the zigzag nanotube. The spectrum of this operator consists of an absolutely continuous part (intervals separated by gaps) plus an infinite number of eigenvalues with infinite multiplicity. They describe all eigenfunctions with the same eigenvalue. They define a Lyapunov function, which is analytic on some Riemann surface. On each sheet, the Lyapunov function has the same properties as in the scalar case, but it has branch points (resonances). They prove that all resonances are real and they determine the asymptotics of the periodic and anti-periodic spectrum and of the resonances at high energy. They show that there exist two types of gaps: i) stable gaps, where the endpoints are periodic and anti-periodic eigenvalues, ii) unstable (resonance) gaps, where the endpoints are resonances (i.e., real branch points of the Lyapunov function). They describe all finite gap potentials. They show that the mapping: potential → all eigenvalues is a real analytic isomorphism for some class of potentials.

Moreover, Korotyaev and Lobanov [KL1] consider magnetic Schrödinger operators on zigzag nanotubes. They describe how the spectrum depends on the magnetic field.

Korotyaev [K2] considers integrated density of states and effective masses for zigzag nanotubes in magnetic fields. He obtains a priori estimates of gap lengths in terms of effective masses.

Kuchment and Post [KuP] consider the case of the zigzag, armchair and achiral nanotubes with $q \in L^2_{even}(0,1)$. They show that the spectrum of the Schrödinger operator (on these nanotubes), as a set, coincides with the spectrum of the Hill operator. We would like to note that the case of even potential q is simple and is closed to the case $q = 0$. Indeed, if q is even, then $F_- = 0$ [MW] and by Theorem 1.3, the corresponding Lyapunov functions are expressed in terms of the Lyapunov function F (for the Hill operator). Moreover, using the identity $F(\lambda) = \cos\kappa(\lambda)$ ($\kappa(\lambda)$ is the quasimomentum for the Hill operator), the Lyapunov function is expressed in terms of $\cos\kappa(\lambda)$, which is similar to the case $q = 0$, where $F(\lambda) = \cos\sqrt{\lambda}$.

In contrast to [KuP] we provide spectral analysis for the Schrödinger operators with arbitrary potentials on edges, that can not be obtained using methods of [KuP].

We present the plan of the paper. In Sect. 2 we prove Theorem 1.2 about the eigenfunctions of H_k and Theorem 1.3 about $\sigma_{ac}(H_k)$. A technical proof of Theorem 1.1 is placed to Sect. 3.

2 Spectrum of the operators H_k

Proof of Theorem 1.2. Proof of i) repeats the arguments from [KL].
ii) Recall $\varkappa_1 = 1 - s^k\varphi_1'^2$, $\varkappa_2 = 1 - s^k\varphi_1'^4$ and $\varphi_1' \neq 0$.

If $k \notin \{0, \frac{N}{2}\}$, then $\operatorname{Im} s^k \neq 0$ which gives $\operatorname{Im}\varkappa_1 \neq 0, \operatorname{Im}\varkappa_2 \neq 0$.

If $k = \frac{N}{2}$, then $s^k = -1$ and $\varkappa_1 = 1 + \varphi_1'^2 \neq 0, \varkappa_2 = 1 + \varphi_1'^4 \neq 0$.

If $k = 0$, then $s^k = 1$ and we have $\varkappa_1 = 1 - \varphi_1'^2, \varkappa_2 = 1 - \varphi_1'^4$. Thus, we obtain
(a) If $\varphi_1'^2 \neq 1$ or $k \neq 0$, then $\varkappa_1 \neq 0, \varkappa_2 \neq 0$.
(b) If $\varphi_1'^2 = 1$ and $k = 0$, then $\varkappa_1 = \varkappa_2 = 0$.
We will define the eigenfunctions of H_k: in the case (a) $\psi^{(0,1)}$ is given by

$$\psi^{(0,1)}_{n,j} = 0,\ \textit{for all}\ n \neq 0,1,\ j \in \mathbb{N}_6,\ \ \textit{and}\ \psi^{(0,1)}_{1,j} = 0,\ j = 5,6, \quad \psi^{(0,1)}_{0,5} = 0, \quad \psi^{(0,1)}_{0,6} = \varkappa_2\varphi,$$

$$s^k\varphi_1'\psi^{(0,1)}_{1,4} = \psi^{(0,1)}_{1,1} = \frac{\psi^{(0,1)}_{1,2}}{\varphi_1'} = \frac{\psi^{(0,1)}_{1,3}}{\varphi_1'^2} = -s^k\varphi_1'\psi^{(0,1)}_{0,4} = -\psi^{(0,1)}_{0,1} = -\frac{\psi^{(0,1)}_{0,2}}{\varphi_1'} = -s^k\varphi_1'^2\psi^{(0,1)}_{0,3} = s^k\varphi_1'^2\varphi, \tag{2.1}$$

and the function $\psi^{(0,2)}$ is given by

$$\psi^{(0,2)}_{n,j} = 0,\ \textit{for all}\ (n,j) \in (\mathbb{Z}\setminus\{-1,0,1\}) \times \mathbb{N}_6,\ \ \textit{and}\ \psi^{(0,2)}_{-1,j} = 0,\ j = 1,3,4,$$

$$\psi^{(0,2)}_{1,j} = 0,\ j \neq 4, \quad \psi^{(0,2)}_{0,1} = 0, \quad \psi^{(0,2)}_{0,3} = \varkappa_1\varphi,$$

$$\psi^{(0,2)}_{-1,2} = -\psi^{(0,2)}_{-1,5} = s^k\psi^{(0,2)}_{0,4} = \frac{\psi^{(0,2)}_{-1,6}}{\varphi_1'} = -\psi^{(0,2)}_{0,2} = \psi^{(0,2)}_{0,5} = -s^k\psi^{(0,2)}_{1,4} = -\varphi_1's^k\psi^{(0,2)}_{0,6} = s^k\varphi_1'\varphi. \tag{2.2}$$

In the case (b) the functions $\psi^{(0,\nu)}, \nu = 1,2$ are given by

$$\psi^{(0,\nu)}_{n,j} = 0,\ (n,j) \in (\mathbb{Z}\setminus\{0,1\}) \times \mathbb{N}_6, \quad \psi^{(0,\nu)}_{1,j} = 0,\ j \in \mathbb{N}_6\setminus\{4\},$$

$$\psi^{(0,\nu)}_{1,4} = \varphi, \quad \psi^{(0,\nu)}_{0,5} = -\varphi, \quad \psi^{(0,\nu)}_{0,6} = \varphi_1'\varphi, \quad \psi^{(0,1)}_{0,1} = \psi^{(0,1)}_{0,3} = -\varphi_1'\varphi,$$

$$\psi^{(0,1)}_{0,2} = 0, \quad \psi^{(0,1)}_{0,4} = -\varphi, \quad \psi^{(0,2)}_{0,1} = \psi^{(0,2)}_{0,3} = \psi^{(0,2)}_{0,4} = 0, \quad \psi^{(0,2)}_{0,2} = \varphi. \tag{2.3}$$

Using (2)-(2), we deduce that $\psi^{(0,\nu)}$ satisfy the Kirchhoff conditions (1.4), (1.5). Thus $\psi^{(0,r)}$ are eigenfunctions of H_k. The operator H_k is periodic, then each $\psi^{(n,\nu)}, (n,\nu) \in \mathbb{Z} \times \mathbb{N}_2$ is an eigenfunction. We will show that the sequence $\psi^{(n,\nu)}, (n,\nu) \in \mathbb{Z} \times \mathbb{N}_2$ forms a basis for $\mathcal{H}_k(\lambda)$. The functions $\psi^{(n,\nu)}$ are linearly independent:
in the case (a) we have $\Gamma_{n,6} \subset \operatorname{supp} \psi^{(n,1)} \setminus \operatorname{supp} \psi^{(m,1)}$ and $\Gamma_{n,3} \subset \operatorname{supp} \psi^{(n,2)} \setminus \operatorname{supp} \psi^{(m,2)}$ for all $n \neq m$ and $\Gamma_{n,1} \subset \operatorname{supp} \psi^{(n,1)} \setminus \operatorname{supp} \psi^{(n,2)}$ for all $n \in \mathbb{Z}$;
in the case (b) we have $\cup_{j \in \mathbb{N}_6, j \neq 4} \Gamma_{n,j} \subset \operatorname{supp} \psi^{(n,\nu)} \setminus \operatorname{supp} \psi^{(m,\nu)}, \nu = 1, 2$, for all $n \neq m$, and $\Gamma_{n,2} \subset \operatorname{supp} \psi^{(n,2)} \setminus \operatorname{supp} \psi^{(n,1)}$ for all $n \in \mathbb{Z}$.

Consider the case (a). For any $f \in \mathcal{H}_k(\lambda)$ we will show the identity (1.12), i.e.,

$$f = \widehat{f}, \quad \textit{where } \widehat{f} = \sum_{(n,\nu) \in \mathbb{Z} \times \mathbb{N}_2} \widehat{f}_{n,\nu} \psi^{(n,\nu)}, \quad \text{and} \quad \widehat{f}_{n,1}, \widehat{f}_{n,2} \quad \text{are given by (1.13).} \tag{2.4}$$

From $\lambda \in \sigma_D$, we deduce that $f|_{\Gamma_{n,j}} = f'_{n,j}(0)\varphi$. The identity $f = \sum \widehat{f}_{n,\nu} \psi^{(n,\nu)}$ and (2), (2) provide

$$\widehat{f}|_{\Gamma_{n,6}} = \widehat{f}_{n,1} \psi^{(n,1)}|_{\Gamma_{n,6}} + \widehat{f}_{n,2} \psi^{(n,2)}|_{\Gamma_{n,6}} = (\widehat{f}_{n,1} \varkappa_2 - \widehat{f}_{n,2})\varphi,$$
$$\widehat{f}|_{\Gamma_{n,3}} = \widehat{f}_{n,1} \psi^{(n,1)}|_{\Gamma_{n,3}} + \widehat{f}_{n,2} \psi^{(n,2)}|_{\Gamma_{n,3}} = (-\widehat{f}_{n,1} + \widehat{f}_{n,2} \varkappa_1)\varphi.$$

Substituting (1.13) into these identities we obtain

$$\widehat{f}|_{\Gamma_{n,6}} = f'_{n,6}(0)\varphi = f|_{\Gamma_{n,6}}, \quad \widehat{f}|_{\Gamma_{n,3}} = f'_{n,3}(0)\varphi = f|_{\Gamma_{n,3}} \quad \textit{all} \quad n \in \mathbb{Z}. \tag{2.5}$$

This yields $\sum |\widehat{f}_{n,\nu}|^2 < \infty$ and $\widehat{f} \in L^2(\Gamma^{(1)})$.

Note that $\widehat{f}$ satisfies the Kirchhoff conditions (1.4), (1.5) and $-\widehat{f}''_\alpha + q\widehat{f}_\alpha = \lambda \widehat{f}_\alpha, \alpha \in \mathcal{Z}_1$. Consider the function $u = f - \widehat{f}$. The function $u = 0$ at all vertices of Γ^1 and then $u_{n,j} = C_{n,j}\varphi, (n,j) \in \mathbb{Z} \times \mathbb{N}_6$, $C_{n,3} = C_{n,6} = 0$. The Kirchhoff boundary conditions (1.4)-(1.5) yield $C_{n,2} = C_{n,4} = 0, n \in \mathbb{Z}$. Assume that $C_{n,1} = C$. Then $C_{n,5} = \frac{C_{n,1}}{\varphi'_1} = \frac{C}{\varphi'_1}$ and $C_{n+1,1} = \frac{C_{n,5}}{\varphi'_1} = \frac{C}{\varphi'^n_1}$ for all $n \in \mathbb{Z}$. Since $u \in L^2(\Gamma^1)$, we have $C = 0$ and $u = 0$, which yields (2.4).

Consider the case (b). For any $f \in \mathcal{H}_k(\lambda)$ we will show the identity (1.12), i.e.,

$$f = \widehat{f}, \quad \textit{where} \quad \widehat{f} = \sum_{n \in \mathbb{Z}, r=1,2} \widehat{f}_{n,r} \psi^{(n,r)}, \qquad \widehat{f}_{n,2} = f'_{n,2}(0), \qquad \widehat{f}_{n,1} = -\varphi'_1 f'_{n,1}(0). \tag{2.6}$$

From (2.6) and $\lambda \in \sigma_D$ we deduce that

$$\widehat{f}|_{\Gamma_{n,1}} = -\widehat{f}_{n,1} \varphi'_1 \varphi = f|_{\Gamma_{n,1}}, \quad \widehat{f}|_{\Gamma_{n,2}} = \widehat{f}_{n,2} \varphi = f|_{\Gamma_{n,2}} \quad \textit{all} \quad n \in \mathbb{Z}. \tag{2.7}$$

This yields $\sum |\widehat{f}_{n,\nu}|^2 < \infty$ and then $\widehat{f} \in \mathcal{H}(\lambda)$.

Consider the function $u = f - \widehat{f}$. The function $u = 0$ at all vertices of Γ^1, which yields $u_{n,j} = C_{n,j}\varphi, n \in \mathbb{Z}, j \in \mathbb{Z}_6$, $C_{n,1} = C_{n,2} = 0$. The Kirchhoff boundary conditions (1.4)-(1.5) yield $C_{n,5} = 0$, and then $C_{n,4} = 0, n \in \mathbb{Z}$. Assume that $C_{n,3} = C$. Then $C_{n,6} = -C_{0,3} = -C$

and $C_{n+1,3} = -C_{n,6} = C$ for all $n \in \mathbb{Z}$. Due to $u \in L^2(\Gamma^{(1)})$, we have $C = 0$ and $u = 0$, which yields (2.6).

The mapping $f \to (\widehat{f}_{n,\nu})_{(n,\nu)\in\mathbb{Z}\times\mathbb{N}_2}$ is a linear and one-to-one mapping from $\mathcal{H}_k$ onto $\ell^2 \oplus \ell^2$. Then it is a linear isomorphism. ■

Proof of Theorem 1.3. (i) Using the standard arguments from [KL] for $D_k(\cdot, \tau) = \det(\mathcal{M}_k - \tau I_4)$ (see also [BBK]) and (1.9), (1.10), (1.11) we obtain (1.15) and (1.14).
The proof of (ii),(iii) repeats standard arguments from [KL].
(iv) If q is even, then $F_- = 0$ (see p.8, [MW]) and identities (1.14) yield

$$F_{0,1} = \frac{(3F+1)^2}{2} - 1, \quad F_{0,2} = \frac{(3F-1)^2}{2} - 1, \tag{2.8}$$

$$F_{k,1} = \frac{1}{2}\biggl(\sqrt{9F^2 - s_k^2} + |c_k|\biggr)^2 - 1, \quad F_{k,2} = \frac{1}{2}\biggl(\sqrt{9F^2 - s_k^2} - |c_k|\biggr)^2 - 1. \tag{2.9}$$

Identities (2.8) show that $\{\lambda \in \mathbb{R} : F_{0,\nu}(\lambda) \in [-1,1] \text{ for some } \nu \in \mathbb{N}_2\} = \{\lambda \in \mathbb{R} : F(\lambda) \in [-1,1]\}$, which together with (1.16) yields $\sigma_{ac}(H_0) = \sigma(\widetilde{H})$. Identities (1.16), (2.9) provide

$$\sigma_{ac}(H_k) = \{\lambda \in \mathbb{R} : F_{k,\nu}(\lambda) \in [-1,1] \text{ for some } \nu \in \mathbb{N}_2\} \subset$$

$$\subset \{\lambda \in \mathbb{R} : F_{0,\nu}(\lambda) \in [-1,1] \text{ for some } \nu \in \mathbb{N}_2\} = \sigma_{ac}(H_0)$$

for all $k \in \mathbb{Z}_N$, which yields $\sigma_{ac}(H_0) = \sigma_{ac}(H)$. ■

Proof of Proposition 1.4. If $q_\varepsilon = \frac{1}{\varepsilon}\delta(t-a), \varepsilon \neq 0, a \in (0,1)$, then we have

$$\vartheta(t,\lambda,q_\varepsilon) = \cos zt + \cos za \frac{\sin z(t-a)}{\varepsilon z}, \quad \varphi(t,\lambda,q_\varepsilon) = \frac{\sin zt}{z} + \frac{\sin za}{\varepsilon z}\frac{\sin z(t-a)}{z},$$

$$F(\lambda,q_\varepsilon) = \cos z + \frac{\sin z}{2z\varepsilon}, \quad F_-(\lambda,q_\varepsilon) = \frac{\sin z(2a-1)}{2z\varepsilon}, \quad z = \sqrt{\lambda}.$$

Let $a = \frac{1}{2} + c_k\varepsilon + \varepsilon^2, k \notin \{0, \frac{N}{2}\}$. Then we obtain

$$F_-(\lambda,q_\varepsilon) = \frac{\sin 2z(c_k\varepsilon + \varepsilon^2)}{2z\varepsilon} = c_k + \varepsilon + O(\varepsilon^2), \quad F_-^2(\lambda,q_\varepsilon) - c_k^2 = 2c_k\varepsilon + O(\varepsilon^2) \tag{2.10}$$

as $|\varepsilon| \to 0$, uniformly on $|z| \leqslant \pi n_0$. Using the identities (1.14), we rewrite the equation $\rho_k(\lambda,q_\varepsilon) = 0, \lambda \in \mathbb{C}$ in the form $F_-^2(\lambda,q_\varepsilon) - c_k^2 = -\frac{9F^2(\lambda,q_\varepsilon)c_k^2}{s_k^2}$, and finally

$$\Phi_+(\lambda,\varepsilon)\Phi_-(\lambda,\varepsilon) = 0, \quad \Phi_\pm(\lambda,\varepsilon) = \varepsilon\biggl(\frac{3F(\lambda,q_\varepsilon)c_k}{s_k} \pm i\sqrt{F_-^2(\lambda,q_\varepsilon) - c_k^2}\biggr), \qquad \lambda \in \mathbb{C}. \tag{2.11}$$

Using $\Phi_\pm(\lambda,0) = \frac{c_k \sin z}{2s_k z}$ and $\frac{\partial\Phi_\pm}{\partial\lambda}((\pi n)^2, 0) \neq 0, n \geqslant 1$, and applying the Implicit Function Theorem to the functions $\Phi_\pm$ we conclude that for each $n \leqslant n_0$ there exists $\varepsilon_n > 0$ such that equation (2.11) has exactly two roots $r_n^\pm(\varepsilon), r_n^\pm(0) = (\pi n)^2$, in the disc $|\varepsilon| < \varepsilon_n$.

Thus we have $\sqrt{r_n^\pm(\varepsilon)} = \pi n + \tau_n^\pm, \tau = \tau_n^\pm \to 0$ as $|\varepsilon| \to 0$ and then

$$F(r_n^\pm(\varepsilon),q_\varepsilon) = (-1)^n\biggl(\cos\tau + \frac{\sin\tau}{2(\pi n + \tau)\varepsilon}\biggr) = \frac{(-1)^n}{2\pi n\varepsilon}(\varepsilon + \tau + O(\tau^2)). \tag{2.12}$$

Substituting (2.10), (2.12) into $F_-^2(\lambda, q_\varepsilon) - c_k^2 = -\frac{9F^2(\lambda,q_\varepsilon)c_k^2}{s_k^2}$, we obtain

$$2c_k\varepsilon + O(\varepsilon^2) = -\frac{9c_k^2}{s_k^2}\left(\frac{\varepsilon + \tau + O(\tau^2)}{2\pi n\varepsilon}\right)^2, \quad \text{and then} \quad \tau = O(\varepsilon). \tag{2.13}$$

Using (2.13) again, we have $\tau = -\varepsilon \pm \frac{2\pi n\varepsilon^{3/2}s_k}{3c_k}\sqrt{-2c_k} + O(\varepsilon^2)$, which yields (1.17). ■

3 Fundamental solutions

Proof of Theorem 1.1. i) Define the operator $\mathcal{S}$ in $\mathbb{C}^N$ by $\mathcal{S}u = (u_N, u_1, \ldots, u_{N-1})^\top$, $u = (u_n)_1^N \in \mathbb{C}^N$. The unitary operator $\mathcal{S}$ has the form $\mathcal{S} = \sum_1^N s^k\mathcal{P}_k$, where $\mathcal{S}e_k = s^k e_k$ and $e_k = \frac{1}{N^{\frac{1}{2}}}(1, s^{-k}, s^{-2k}, ..., s^{-kN+k})$ is an eigenvector (recall $s = e^{i\frac{2\pi}{N}}$); $\mathcal{P}_k u = e_k(u, e_k)$ is a projector. The function f in the Kirchhoff boundary conditions (1.2) is a vector function $f = (f_\omega), \omega = (n, j, k) \in \mathcal{Z}$. We define a new vector-valued function $f_{n,j} = (f_{n,j,k})_{k=1}^N \in \mathbb{C}^N, (n, j) \in \mathcal{Z}_1 = \mathbb{Z} \times \mathbb{N}_6$, which satisfies the equation $-f_{n,j}'' + qf_{n,j} = \lambda f_{n,j}$, and the conditions for all $n \in Z$, which follow from the Kirchhoff conditions (1.2):

$$\begin{aligned} &f_{n,1}(1) = f_{n,2}(0) = f_{n,5}(0), \quad f_{n,2}(1) = f_{n,3}(0) = f_{n,6}(0), \\ &\qquad f_{n,3}(1) = f_{n,4}(0) = f_{n-1,6}(1), \quad \mathcal{S}f_{n,4}(1) = f_{n,1}(0) = f_{n-1,5}(1), \end{aligned} \tag{3.1}$$

$$\begin{aligned} &f_{n,1}'(1) - f_{n,2}'(0) - f_{n,5}'(0) = 0, \quad f_{n,2}'(1) - f_{n,3}'(0) - f_{n,6}'(0) = 0, \\ &\qquad f_{n,3}'(1) - f_{n,4}'(0) + f_{n-1,6}'(1) = 0, \quad \mathcal{S}f_{n,4}'(1) - f_{n,1}'(0) + f_{n-1,5}'(1) = 0. \end{aligned} \tag{3.2}$$

The operators $\mathcal{S}$ and $\mathscr{H}$ commute, then we deduce that $\mathscr{H}\mathcal{P}_k$ is unitarily equivalent to the operator H_k acting in $L^2(\Gamma^1)$ and H_k is given by $(H_k f)_\alpha = -f_\alpha'' + q(t)f_\alpha$, where $(f_\alpha)_{\alpha\in\mathcal{Z}_1}, (f_\alpha'')_{\alpha\in\mathcal{Z}_1} \in L^2(\Gamma^1)$ and components f_α satisfy the boundary conditions (1.4), (1.5). Thus $\mathscr{H}$ is unitarily equivalent to the operator $H = \oplus_1^N H_k$

ii) We prove the following idenities:

$$\mathcal{M}_k = \mathcal{R}^{-1}Y\mathcal{T}_k\mathcal{R}, \quad \mathcal{T}_k = \begin{pmatrix} V_k & I_2 \\ V_0V_k - I_2 & V_0 \end{pmatrix}, \quad V_k = \begin{pmatrix} 2F & -s^k \\ -s^{-k} & 2F \end{pmatrix}, \quad Y = \begin{pmatrix} \vartheta_1 I_2 & I_2 \\ \varphi_1\vartheta_1' I_2 & \varphi_1' I_2 \end{pmatrix}. \tag{3.3}$$

Let $\varphi_t = \varphi(t, \lambda), \varphi_t' = \varphi'(t, \lambda), ...$ Recall that any solution y of the equation $-y'' + qy = \lambda y$ on $[0, 1]$ satisfies

$$y(t) = y(0)\vartheta_t + \frac{\varphi_t}{\varphi_1}(y(1) - \vartheta_1 y(0)), \quad \begin{pmatrix} y(1) \\ y'(1) \end{pmatrix} = \mathcal{M}\begin{pmatrix} y(0) \\ y'(0) \end{pmatrix}, \quad \mathcal{M}^{-1} = \begin{pmatrix} \varphi_1' & -\varphi_1 \\ -\vartheta_1' & \vartheta_1 \end{pmatrix}. \tag{3.4}$$

Let $\mathcal{F}_n(t) = (f_{n,5}(t), f_{n,6}(t), f_{n,5}'(t), f_{n,6}'(t))^\top$. We will prove that for each vector $\mathcal{F}_{-1}(1) = h = (h_5, h_6, h_5', h_6')^\top \in \mathbb{C}^4$ there exists a unique vector function $(f_\alpha)_{\alpha\in\mathcal{Z}_1}$, satisfying the equations $-f_\alpha'' + q(x)f_\alpha = \lambda f_\alpha, x \in [0, 1], \alpha \in \mathcal{Z}_1$, and the Kirchhoff boundary conditions (1.4), (1.5).

Firstly, we will determine $f_{0,1}, f_{0,3}$ in terms of h. The Kirchhof conditions (1.4), (1.5) at $n = 0$ yield

$$f_{0,3}(1) = f_{0,4}(0) = h_6, \quad s^k f_{0,4}(1) = f_{0,1}(0) = h_5,$$
$$f'_{0,3}(1) - f'_{0,4}(0) + h'_6 = 0, \quad s^k f'_{0,4}(1) - f'_{0,1}(0) + h'_5 = 0. \tag{3.5}$$

Let $w_{13}(t) = (f_{0,1}(t), f_{0,3}(1-t), f'_{0,1}(t), f'_{0,3}(1-t))^\top$. Then conditions (3.5) imply

$$w_{13}(0) = \begin{pmatrix} I_2 & 0 \\ 0 & \mathbf{j}_1 \end{pmatrix} h + w_4, \quad w_4 = (0, 0, s^k f'_{0,4}(1), f'_{0,4}(0))^\top, \quad \mathbf{j}_1 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \tag{3.6}$$

Using (3.4), (3.5) we get $f_{0,4}(t) = h_6 \vartheta_t + \frac{\varphi_t}{\varphi_1}(s^{-k} h_5 - \vartheta_1 h_6), \quad t \in [0,1]$, which yields

$$f'_{0,4}(0) = \frac{1}{\varphi_1}(s^{-k} h_5 - \vartheta_1 h_6), \quad f'_{0,4}(1) = h_6 \vartheta'_1 + \frac{\varphi'_1}{\varphi_1}(s^{-k} h_5 - \vartheta_1 h_6) = \frac{1}{\varphi_1}(s^{-k} \varphi'_1 h_5 - h_6),$$

where we have used the identity $\vartheta_1 \varphi'_1 - \vartheta'_1 \varphi_1 = 1$. The last identities and (3.6) give

$$w_4 = \begin{pmatrix} 0 & 0 \\ \frac{A_0}{\varphi_1} & 0 \end{pmatrix} h, \quad A_0 = \begin{pmatrix} \varphi'_1 & -s^k \\ s^{-k} & -\vartheta_1 \end{pmatrix}, \qquad w_{13}(0) = Y_0 h, \quad Y_0 = \begin{pmatrix} I_2 & 0 \\ \frac{A_0}{\varphi_1} & \mathbf{j}_1 \end{pmatrix}. \tag{3.7}$$

Thus for each $h = \mathcal{F}_{-1}(1) \in \mathbb{C}^4$ there exists unique $w_{13}(0) = (f_{0,1}(0), f_{0,3}(1), f'_{0,1}(0), f'_{0,3}(1))^\top$ and then the unique functions $f_{0,1}(t), f_{0,3}(t), t \in [0,1]$ satisfying the equation $-f'' + qf = \lambda f$ and the Kirchhoff boundary conditions (3.5).

Identities (3.4) imply $\begin{pmatrix} f_{0,1}(1) \\ f'_{0,1}(1) \end{pmatrix} = \mathcal{M} \begin{pmatrix} f_{0,1}(0) \\ f'_{0,1}(0) \end{pmatrix}, \quad \begin{pmatrix} f_{0,3}(0) \\ f'_{0,3}(0) \end{pmatrix} = \mathcal{M}^{-1} \begin{pmatrix} f_{0,3}(1) \\ f'_{0,3}(1) \end{pmatrix}$, which yields

$$w_{13}(1) = Y_1 w_{13}(0), \qquad Y_1 = \begin{pmatrix} A_1 \mathbf{j}_2 & \varphi_1 \mathbf{j}_1 \\ \vartheta'_1 \mathbf{j}_1 & \mathbf{j}_2 A_1 \end{pmatrix}, \quad A_1 = \begin{pmatrix} 0 & \vartheta_1 \\ \varphi'_1 & 0 \end{pmatrix}, \quad \mathbf{j}_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \tag{3.8}$$

Secondly, we will determine $f_{0,5}, f_{0,6}$ in terms of $f_{0,1}, f_{0,3}$. The Kirchhof conditions (1.4), (1.5) at $n = 0$ yield

$$f_{0,1}(1) = f_{0,2}(0) = f_{0,5}(0), \quad f_{0,2}(1) = f_{0,3}(0) = f_{0,6}(0),$$
$$f'_{0,1}(1) - f'_{0,2}(0) - f'_{0,5}(0) = 0, \quad f'_{0,2}(1) - f'_{0,3}(0) - f'_{0,6}(0) = 0, \tag{3.9}$$

which gives

$$\mathcal{F}_0(0) = \begin{pmatrix} I_2 & 0 \\ 0 & \mathbf{j}_1 \end{pmatrix} w_{13}(1) + w_2, \quad w_2 = (0, 0, -f'_{0,2}(0), f'_{0,2}(1))^\top, \tag{3.10}$$

where $w_{13}(1) = (f_{0,1}(1), f_{0,3}(0), f'_{0,1}(1), f'_{0,3}(0))^\top$, $\mathcal{F}_0(0) = (f_{0,5}(0), f_{0,6}(0), f'_{0,5}(0), f'_{0,6}(0))^\top$. Using the Kirchhof conditions (3) and (3.4), we obtain

$$f_{0,2}(t) = f_{0,1}(1) \vartheta_t + \frac{\varphi_t}{\varphi_1}(f_{0,3}(0) - \vartheta_1 f_{0,1}(1)), \quad f'_{0,2}(0) = \frac{1}{\varphi_1}(f_{0,3}(0) - \vartheta_1 f_{0,1}(1)),$$

$$f_{0,2}'(1) = f_{0,1}(1)\vartheta_1' + \frac{\varphi_1'}{\varphi_1}(f_{0,3}(0) - \vartheta_1 f_{0,1}(1)) = \frac{1}{\varphi_1}(\varphi_1' f_{0,3}(0) - f_{0,1}(1)),$$

where we have used the Wronskian $\vartheta_1\varphi_1' - \vartheta_1'\varphi_1 = 1$. Then

$$w_2 = \begin{pmatrix} 0 & 0 \\ \frac{1}{\varphi_1}A_2 & 0 \end{pmatrix} w_{13}(1), \qquad A_2 = \begin{pmatrix} \vartheta_1 & -1 \\ -1 & \varphi_1' \end{pmatrix}. \tag{3.11}$$

Substituting this identity into (3.10) we obtain

$$\mathcal{F}_0(0) = Y_2 u(1), \qquad Y_2 = \begin{pmatrix} I_2 & 0 \\ \frac{1}{\varphi_1}A_2 & \mathbf{j}_1 \end{pmatrix}. \tag{3.12}$$

Thus for each $h \in \mathbb{C}^4$ there exists a unique vector $\mathcal{F}_0(0) = (f_{0,5}(0), f_{0,6}(0), f_{0,5}'(0), f_{0,6}'(0))^\top$ and the unique functions $f_{0,5}(t), f_{0,6}(t), t \in [0,1]$ satisfying the equation $-f'' + qf = \lambda f$ and the Kirchhoff boundary conditions (3). Identities (3.4) give $\begin{pmatrix} f_{0,5}(1) \\ f_{0,5}'(1) \end{pmatrix} = \mathcal{M}\begin{pmatrix} f_{0,5}(0) \\ f_{0,5}'(0) \end{pmatrix}$ and $\begin{pmatrix} f_{0,6}(1) \\ f_{0,6}'(1) \end{pmatrix} = \mathcal{M}\begin{pmatrix} f_{0,6}(0) \\ f_{0,6}'(0) \end{pmatrix}$, then

$$\mathcal{F}_0(1) = Y_3\mathcal{F}_0(0), \qquad Y_3 = \begin{pmatrix} \vartheta_1 I_2 & \varphi_1 I_2 \\ \vartheta_1' I_2 & \varphi_1' I_2 \end{pmatrix}. \tag{3.13}$$

Thus we have proved that for each $h \in \mathbb{C}^4$ there exist the unique functions $f_{0,j}(t), j \in \mathbb{N}_6$, satisfying the equation $-f'' + qf = \lambda f$ and the Kirchhoff boundary conditions (1.4), (1.5). By the periodicity, we obtain similar results for all functions $f_\alpha, \alpha \in \mathcal{Z}_1$.

Recall that the fundamental solutions $\vartheta_k^{(\nu)} = (\vartheta_{k,\alpha}^{(\nu)})_{\alpha\in\mathcal{Z}_1}, \varphi_k^{(\nu)} = (\varphi_{k,\alpha}^{(\nu)})_{\alpha\in\mathcal{Z}_1}, \nu = 1,2$ satisfy (1.6) and the monodromy matrix $\mathcal{M}_k$ is given by (1.8). Then $\mathcal{F}_0(1) = \mathcal{M}_k h,\ h = \mathcal{F}_{-1}(1)$, and identities (3.7), (3.8), (3.12), (3.13) yield

$$\mathcal{M}_k = Y_3 Y_2 Y_1 Y_0. \tag{3.14}$$

Now we calculate $Y_2Y_1Y_0$. We need the identities

$$\varphi_1' + \vartheta_1 = 2F, \qquad \vartheta_1'\varphi_1 + \varphi_1'^2 = 2\varphi_1' F - 1, \qquad \vartheta_1'\varphi_1 + \vartheta_1^2 = 2\vartheta_1 F - 1. \tag{3.15}$$

Identities (3.7), (3.8) give

$$Y_1Y_0 = \begin{pmatrix} A_1\mathbf{j}_2 + \mathbf{j}_1 A_0 & \varphi_1 I_2 \\ \vartheta_1'\mathbf{j}_1 + \frac{1}{\varphi_1}\mathbf{j}_2 A_1 A_0 & \mathbf{j}_2 A_1 \mathbf{j}_1 \end{pmatrix} = \begin{pmatrix} V_k & \varphi_1 I_2 \\ \frac{1}{\varphi_1}B_1 & B_2 \end{pmatrix}, \qquad B_2 = \begin{pmatrix} \varphi_1' & 0 \\ 0 & -\vartheta_1 \end{pmatrix}, \tag{3.16}$$

$$B_1 = \begin{pmatrix} \varphi_1\vartheta_1' & 0 \\ 0 & -\varphi_1\vartheta_1' \end{pmatrix} + \begin{pmatrix} \varphi_1' & 0 \\ 0 & \vartheta_1 \end{pmatrix}\begin{pmatrix} \varphi_1' & -s^k \\ s^{-k} & -\vartheta_1 \end{pmatrix} = \begin{pmatrix} 2\varphi_1' F - 1 & -s^k\varphi_1' \\ s^{-k}\vartheta_1 & 1 - 2\vartheta_1 F \end{pmatrix}, \tag{3.17}$$

where we have used (3.15) and V_k are given by (3.3). Identities (3.12), (3.16) imply

$$Y_2Y_1Y_0 = \begin{pmatrix} I_2 & 0 \\ \frac{1}{\varphi_1}A_2 & \mathbf{j}_1 \end{pmatrix}\begin{pmatrix} V_k & \varphi_1 I_2 \\ \frac{1}{\varphi_1}B_1 & B_2 \end{pmatrix} = \begin{pmatrix} V_k & \varphi_1 I_2 \\ \frac{1}{\varphi_1}(A_2 V_k + \mathbf{j}_1 B_1) & A_2 + \mathbf{j}_1 B_2 \end{pmatrix} = \mathcal{R}^{-1}\mathcal{T}_k\mathcal{R}, \tag{3.18}$$

where

$$\mathcal{T}_k = \begin{pmatrix} V_k & I_2 \\ A_2V_k + \mathbf{j}_1 B_1 & A_2 + \mathbf{j}_1 B_2 \end{pmatrix}. \tag{3.19}$$

Using (3.11), (3.16), (3.17) we have

$$A_2V_k + \mathbf{j}_1B_1 = \begin{pmatrix} \vartheta_1 & -1 \\ -1 & \varphi_1' \end{pmatrix}\begin{pmatrix} 2F & -s^k \\ -s^{-k} & 2F \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} 2\varphi_1'F - 1 & -s^k\varphi_1' \\ s^{-k}\vartheta_1 & -2\vartheta_1F + 1 \end{pmatrix} = V_0V_k - I_2,$$

$$A_2 + \mathbf{j}_1B_2 = \begin{pmatrix} \vartheta_1 & -1 \\ -1 & \varphi_1' \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} \varphi_1' & 0 \\ 0 & -\vartheta_1 \end{pmatrix} = V_0.$$

Substituting the last two identities into (3.19), we obtain the identity for $\mathcal{T}_k$ in (3.3). Substituting identity (3.18) into (3.14), we obtain $\mathcal{M}_k = Y_3\mathcal{R}^{-1}\mathcal{T}_k\mathcal{R}$ which yields (3.3). Identities (3.3) show that the matrix-valued function $\mathcal{R}\mathcal{M}_k\mathcal{R}^{-1}$ is entire.

Identities (3.7), (3.12) give $\det Y_0 = \det Y_2 = -1$. Identity (3.8) shows that Y_1 is obtained from the matrix $\begin{pmatrix} \mathcal{M} & 0 \\ 0 & \mathcal{M} \end{pmatrix}$ by the transposition of rows and columns, hence $\det Y_1 = 1$. (3.13) implies that Y_1 is obtained from the matrix $\begin{pmatrix} \mathcal{M} & 0 \\ 0 & \mathcal{M}^{-1} \end{pmatrix}$ by the transposition of rows and columns, hence $\det Y_3 = \det \mathcal{M} \det \mathcal{M}^{-1} = 1$. Then (3.14) gives $\det \mathcal{M}_k = 1$.

Furthermore, (3.3) gives

$$\operatorname{Tr} \mathcal{M}_k = \operatorname{Tr} \begin{pmatrix} \vartheta_1 I_2 & \varphi_1 I_2 \\ \vartheta_1' I_2 & \varphi_1' I_2 \end{pmatrix}\begin{pmatrix} V_k & \varphi_1 I_2 \\ \frac{1}{\varphi_1}(V_0V_k - I_2) & V_0 \end{pmatrix} = \operatorname{Tr}(\vartheta_1 V_k + V_0V_k - I_2 + \varphi_1\vartheta_1' I_2 + \varphi_1' V_0),$$

$$\operatorname{Tr} V_k = \operatorname{Tr} V_0 = 4F, \quad \operatorname{Tr}(V_0V_k) = 8F^2 + s^{-k} + s^k. \tag{3.20}$$

Hence

$$\operatorname{Tr} \mathcal{M}_k = 4F(\vartheta_1 + \varphi_1') + 8F^2 + s^{-k} + s^k - 2 + 2\varphi_1\vartheta_1' = 16F^2 + s^{-k} + s^k - 2 + 2\varphi_1\vartheta_1' = \operatorname{Tr} \mathcal{M}_0 - 4s_k^2,$$

where $\operatorname{Tr} \mathcal{M}_0 = 16F^2 + 2\varphi_1\vartheta_1'$, which yields (1.9).

We prove (1.10). We have

$$\operatorname{Tr} \mathcal{M}_k^2 = \operatorname{Tr}(\vartheta_1V_k + V_0V_k - I_2)^2 + 2\operatorname{Tr}(\vartheta_1 I_2 + V_0)(\varphi_1\vartheta_1'V_k + \varphi_1'(V_0V_k - I_2)) + \operatorname{Tr}(\varphi_1\vartheta_1' I_2 + \varphi_1'V_0)^2. \tag{3.21}$$

Direct calculations give

$$\operatorname{Tr} V_k^2 = \operatorname{Tr} V_0^2 = 8F^2 + 2, \quad \operatorname{Tr}(V_0V_k)^2 = 32F^4 + 16F^2(s^k + s^{-k} + 1) + s^{2k} + s^{-2k}, \tag{3.22}$$

$$\operatorname{Tr}(\vartheta_1 V_k^2 V_0 + \varphi_1' V_0^2 V_k) = 32F^4 + 8F^2(s^k + s^{-k} + 1). \tag{3.23}$$

Substituting identities (3.20), (3.22) and (3.23) into (3.21) we obtain

$$\operatorname{Tr} \mathcal{M}_k^2 = 128F^4 + 8F^2 + 32F^2\varphi_1\vartheta_1' + (32F^2 + 4\varphi_1\vartheta_1')(s^k + s^{-k}) + s^{2k} + s^{-2k} - 8\vartheta_1\varphi_1' + 2\varphi_1^2\vartheta_1'^2 + 2$$

$$= \frac{1}{2}(\operatorname{Tr} \mathcal{M}_0)^2 + 72F^2 - 8s_k^2 \operatorname{Tr} \mathcal{M}_0 - 4s_{2k}^2 - 4 = \operatorname{Tr} \mathcal{M}_0^2 - 8s_k^2 \operatorname{Tr} \mathcal{M}_0 - 4s_{2k}^2,$$

where we have used the identities $s^k+s^{-k}=2-4s_k^2, \vartheta_1\varphi_1'=\varphi_1\vartheta_1'+1$. This yields (1.10).

Now we will prove (1.11). Direct calculation show that $Y^\top JY=J$ and

$$\mathcal{T}_k^\top J\mathcal{T}_k=\begin{pmatrix} V_k^\top\mathbf{j}_2(V_0V_k-I_2)-(V_0V_k-I_2)^\top\mathbf{j}_2V_k & V_k^\top\mathbf{j}_2V_0-(V_0V_k-I_2)^\top\mathbf{j}_2 \\ \mathbf{j}_2(V_0V_k-I_2)-V_0\mathbf{j}_2V_k & \mathbf{j}_2V_0-V_0\mathbf{j}_2 \end{pmatrix}=J,$$

where the identities $\mathbf{j}_2V_k=V_k^\top\mathbf{j}_2, \mathbf{j}_2V_0=V_0\mathbf{j}_2$ was used. Then $(Y\mathcal{T}_k)^\top JY\mathcal{T}_k=J$. Using the identities $\mathcal{R}J\mathcal{R}=\varphi_1 J$, $\mathcal{R}^{-1}J\mathcal{R}^{-1}=\frac{1}{\varphi_1}J$, we have (1.11). ■

Acknowledgments. Evgeny Korotyaev was partly supported by DFG project BR691/23-1. The various parts of this paper were written at ESI, Vienna , E. Korotyaev is grateful to the Institute for the hospitality. A. Badanin is grateful to the Mathematical Institute of Humboldt Univ. for the hospitality.

References

[A] Alexander S. Superconductivity of networks. A percolation approach to the effect of disorder. Phys. Rev. B 27 (1983), 1541–1557.

[AL] Anantram M.P., F. Léonard. Physics of carbon nanotube electronic devices. Rep. Prog. Phys. 69 (2006), 507–561.

[AT] Adamyan V., Tishchenko S. One-electron states and interband optical absorption in single-wall carbon nanotubes. J. Phys. Condens. Matter 19 (2007), 621-6218.

[BBK] Badanin, A.; Brüning, J.; Korotyaev, E. The Lyapunov function for Schrödinger operators with a periodic 2×2 matrix potential. J. Funct. Anal. 234 (2006), no. 1, 106–126.

[CK] Chelkak, D.; Korotyaev, E. Spectral estimates for Schrödinger operators with periodic matrix potentials on the real line. Int. Math. Res. Not. 2006, Art. ID 60314, 41 pp.

[C] Coulson C. A. Note on the applicability of the free-electron network model to metals. Proc. Phys. Soc. A 67 (1954), 608–614.

[HSO] Hamada N., Sawada Sh., Oshiyama A. New one-dimensional conductors: graphitic microtubules. Phys. Rev. Lett. 9 (1992), no 10, 1579–1581.

[Ha] Harris P. Carbon Nanotubes and Related Structures, Cambridge Univ. Press., Cambridge, 1999

[HRB] Hod O., Rabani E., Baer R. Magnetoresistance devices based on signle-walled carboon nanotubes. J. Chem. Phys. 123 (2005), no. 051103, 1-4.

[Ii] Iijima S. Helical microtubules of graphitic carbon. Nature 354 (1991), 56–58.

[K1] Korotyaev, E. Conformal spectral theory for the monodromy matrix. Preprint, 2006

[K2] Korotyaev, E. Effective masses for zigzag nanotubes in magnetic fields, preprint 2007.

[KL] Korotyaev, E.; Lobanov, I. Schrödinger operators on zigzag periodic graphs, To be published in Ann. Inst. H.Poincare.

[KL1] Korotyaev, E.; Lobanov, I. Zigzag periodic nanotube in magnetic field, preprint 2006

[Ku] Kuchment, P. Graph models for waves in thin structures, *Waves in Random Media*, **12** (2002), R1-R24

[KuP] Kuchment, P.; Post, O. On the spectra of carbon nano-structures, preprint 2006

[MDW] Mintmire J. W., Dunlap B. I., White C. T. Are fullerene tubules metallic? Phys. Rev. Lett. 68 (1992), no. 2, 631–634.

[M] Montroll E. W. Quantum theory on a network. I. A solvable model whose wavefunctions are elementary functions. J. Math. Phys. 11 (1970), no. 2, 635–648.

[MW] Magnus, W.; Winkler, S. Hill's equation. Dover Publications, Inc., New York, 1979

[MV] Molchanov, S.; Vainberg, B. Slowing down of the wave packets in quantum graphs. Waves Random Complex Media 15(2005), 101–112

[OHKL] Odom T. W., Huang J.-L., Kim P., Lieber C. M., Structure and Electronic Properties of Carbon Nanotubes. J. Phys. Chem. B 104 (2000), no. 13, 2794.

[Pa] Pauling, L. The diamagnetic anisotropy of aromatic molecules, *Journal of Chemical Physics*, 4, 1936, 673-677

[PBRL] Pacheco M., Barticevic Z., Rocha C. G., Latgé A. Electric field effects on the energy spectrum of carbon nanotubes. J. Phys. Condens. Matter 17 (2005), 5839–5847.

[RMT] Reich S., Maultzsch J., Thomas C. Tight-binding description of graphene. Phys. Rev. B 66 (2002), no. 035412. 1–5.

[RS] Ruedenberg, K.; Scherr, C.W. Free-electron network model for conjugated systems. I. Theory, *The Journal of Chemical Physics*, 21, 1953, 1565-1581.

[SDD] Saito, R.; Dresselhaus, G.; Dresselhaus, M. Physical properties of carbon nanotubes, Imperial College Press, 1998

[SFDD] Saito R., Fujita M., Dresselhaus G., Dresselhaus M. S. Electronic structure of graphene tubules based on C_{60}. Phys. Rev. B 46 (1992), no 3, 1804–1811.